

SHORT COMMUNICATION

Sexual differences in the external measurements of Black-tailed Gulls breeding on Rishiri Island, Japan**ORNITHOLOGICAL
SCIENCE**© The Ornithological Society
of Japan 2002Michiyo CHOCHI^{1,*}, Yasuaki NIIZUMA^{2,#,**} and Masaoki TAKAGI³¹ Department of Ecology and Systematics, Graduate School of Agriculture, Hokkaido University, Sapporo 060–8589, Japan² Institute of Low Temperature Science, Hokkaido University, Sapporo 060–0819, Japan³ Laboratory of Animal Sociology, Faculty of Science, Osaka City University, Sumiyoshi-ku, Osaka, 558–8585, Japan

Sex ratio is an important parameter to consider in ecological and conservation seabirds studies (Weimerskirch & Jouventin 1987). Visually assessing with certainty the sex of live seabirds, however, may be an impossible task when no obvious differences in plumage or body size exist between the two sexes. Several previous studies of Laridae species have shown that males are significantly larger than females (external measurements) and discriminant functions using measurements sex Laridae with great reliability (Shugart 1977; Ryder 1978; Fox et al. 1981; Schnell et al. 1985; Coulson et al. 1983; Evans et al. 1993; Bosch 1996; Rodriguez et al. 1996; Palomares et al. 1997).

The Black-tailed Gull *Larus crassirostris* is a medium-sized gull, endemic to the northwest Pacific, breeding extensively in coastal regions around the Japanese archipelago. Although Black-tailed Gulls are one of the most common seabird species in Japan, little is known about their biology or morphology. Similarly to other Laridae species, Black-tailed Gulls have no sexual differences in their plumage or coloration. This study is aimed at describing the external measurements of Black-tailed Gulls and determining a discriminant function using the measurements to facilitate the sexing of the gulls in the field.

METHODS

A management program controlling the number of Black-tailed Gulls was carried out by the Rishiri

town office on Rishiri Island (45°05'N, 141°07'E), off the northwest coast of Hokkaido, Japan. In this region, gulls are a nuisance to townspeople, inflicting damage on the commercial fishery and affecting flights to and from the airport nearby. Carcasses of gulls culled during the course of this program were used to investigate the relationship between the external measurements and sex.

On Rishiri Island, about 7,900 pairs bred in 1998 and 8,600 pairs in 1999 (Hokkaido Government 1999, 2000), on a 9.12 ha area covered with dwarf bamboo *Sasa nipponica*. Gulls were shot around the colonies from 14–18 June 1998 during the incubation period, and from 17–24 June 1999 during the incubation and hatching periods (Hokkaido 1999, 2000). A total of 237 carcasses was collected and frozen less than one hour after being shot. External measurements were carried out after the carcasses had been thawed.

Ten external measurements were performed following the procedures described in Bosch (1996). Head length (HL), long bill length (LBL), short bill length (SBL), nalospi (Nal: the distance from the tip of bill to the nostril), bill depth (BD), tarsus length (Tar), foot length (FL), and middle toe length (MTL) were measured to the nearest 0.01 mm using vernier calipers. Tail length (Tail L: the length between the uropygial gland and the tip of one rectrix) and wing length (WL: natural chord) were measured to the nearest 0.1 mm and 1 mm, respectively, using rulers. Body mass was weighed to the nearest 0.1 g using an electrical balance. Several studies have found sexual dimorphism in body mass for Laridae species (Ingolfsson 1969; Ryder 1978; Threlfall & Jewer 1978; Hunt et al. 1980; Fox et al. 1981; Coulson et al. 1983; Monaghan et al. 1983; Bosch 1996; Rodriguez et al. 1996; Palomares et al. 1997). However, due to its variability throughout the breeding season (Jones

(Received 14 May 2002; Accepted 12 July 2002)

Corresponding author, E-mail: niizuma@affrc.go.jp

* Present address: Yokohama Zoological Gardens, Kamishirane-cho 1175–1, Asahi-ku, Yokohama 241–0001, Japan.

** Present address: Japan Society for the Promotion of Science, Hokkaido National Fisheries Research Institute, Katsurakoi 116, Kushiro 085–0802, Japan.

Table 1. Average values for external measurements and MD index of Black-Tailed Gull. Standard deviations are expressed as \pm SD, and ranges are given in Parentheses. Significance levels were improved using the sequential Bonferroni technique.

Adult	Male (N=139)	Female (N=98)	t-value	P-value	MD index
Head Length (mm)	112.78 \pm 2.66 (105.60–118.99)	103.99 \pm 2.53 (96.82–111.35)	25.55	<0.01	8.11
Long Bill Length (mm)	74.68 \pm 2.84 (62.70–85.20)	69.03 \pm 2.77 (63.10–77.40)	15.24	<0.01	7.86
Short Bill Length (mm)	52.21 \pm 2.12 (46.75–57.10)	47.67 \pm 2.16 (42.40–55.35)	16.11	<0.01	9.09
Nalospa (mm)	24.53 \pm 1.59 (21.44–28.80)	22.39 \pm 1.51 (18.70–28.90)	10.40	<0.01	9.12
Bill Depth (mm)	17.11 \pm 1.24 (14.50–20.67)	15.09 \pm 1.01 (13.30–18.90)	13.32	<0.01	12.55
Tarsus Length (mm)	57.62 \pm 2.15 (49.85–63.35)	53.19 \pm 2.10 (45.60–57.45)	15.76	<0.01	8.00
Foot Length (mm)	104.47 \pm 3.52 (89.45–114.12)	97.36 \pm 3.46 (89.15–111.98)	15.42	<0.01	7.05
Middle Toe Length (mm)	52.12 \pm 2.25 (43.45–57.04)	48.28 \pm 2.34 (39.95–52.01)	12.72	<0.01	7.65
Tail Length (mm)	143.85 \pm 5.83 (126.5–157.0)	136.36 \pm 4.84 (124.3–149.0)	10.43	<0.01	5.35
Wing Length (cm)	39.01 \pm 0.80 (36.5–157.0)	37.39 \pm 0.75 (35.6–39.2)	15.65	<0.01	4.24

1994; Croxall 1995), sexual difference in body mass is not described here (Lorentsen & Røy 1994). Once these measurements were collected, birds were dissected and sexed based on their reproductive organs.

Each external measurement was compared between the sexes using Student's *t*-test. Sequential Bonferroni's *t*-test was performed to increase significance power. To compare the degree of sexual difference among measurements, the mean difference (MD) index was used (Agnew & Kerry 1995). The MD index was expressed as the difference of the average measurements between male and female:

$$MD = 200 \times (X.m - X.f) / (X.m + X.f)$$

where *X.m* and *X.f* are the mean of the measurements for males and females, respectively; a larger index indicating greater sexual dimorphism.

Finally, a discriminant function analysis was performed using HL and BD for sex determination of the gull, because these two measurements are readily taken during field studies (Fox et al. 1981; Palomares et al. 1997).

RESULTS

Male gulls were significantly larger than females for all external measurements (Table 1). MD indices

for parameters of the head region, i.e. HL, LBL, SBL, Nal, and BD were larger than those of the extremities, i.e. WL and Tail L. The average body mass of males was 641.2 g \pm 44.0 (SD) and 537.5 g \pm 42.3 for females.

Using HL and BD measurements, the following discriminant function was obtained;

$$D = 150.63 - 1.22 HL - 1.14 BD \\ (F_{2,234} = 372.88, P < 0.0001)$$

Both two variables contributed significantly to the function (HL, $F_{1,234} = 324.21$, $P < 0.0001$; BD, $F_{1,234} = 25.41$, $P < 0.0001$). Following this function, birds were classified as males when $D < 0$ and female when $D > 0$. This discriminant function proved reliable for 95.7% of the males (133 out of 139), for 98.0% of the females (96 out of 98), and for a total of 96.6% of the birds (Fig. 1).

DISCUSSION

This study demonstrated that Black-tailed Gulls, breeding on Rishiri Island have marked sexual differences in their external measurements and can be sexed by a discriminant function using HL and BD with a total reliability of 96.6%. This discriminant function will prove useful for sexing Black-tailed




Fig. 1. Relationship between bill depth (BD) and head length (HL) to the function $1.22 \text{ HL} = 150.63 - 1.14 \text{ BD}$. Open circles show females and closed circles show males.

Gulls in the field. Rapid sexing of a bird would contribute to shortening handling time thereby allowing researchers to release birds quickly after a minimum of disturbance. Before applying this method to other populations or age classes, however, sexing accuracy requires further testing, especially since Laridae species show inter-colonial variation in external measurements (Threlfall & Jewer 1978; Monaghan et al. 1983; Jehl 1987; Evans et al. 1993), as well as age related differences (Coulson et al. 1981; Allaine & Lebreton 1990; Palomares et al. 1997). The greater difference in size between the sexes found in the head region indicates that this region, especially the bill, may play an important role, probably in sexual display and territorial defence by males (Ingolfsson 1969). The smaller sexual dimorphism in the wing region may be related to the species' flight performance (Schnell et al. 1985; Croxall 1995).

ACKNOWLEDGMENTS

We would like to thank the Rishiri town office for authorizing us to use the gulls' carcasses and for supporting our work on Rishiri Island. We are grateful to M. Sato and M. Senda for assistance in the field and Y. Ropert-Coudert for helpful comments on the manuscript.

REFERENCES

- Agnew DJ & Kerry KR (1995) Sexual dimorphism in penguins. In: Dann P, Norman I & Reilly P (eds) *The Penguins: ecology and management*. pp 299–318. Surrey Beatty and Sons, Australia.
- Allaine D & Lebreton J-D (1990) The influence of age and sex on wing-tip pattern in adult Black-headed Gulls *Larus ridibundus*. *Ibis* 132: 560–567.
- Bosch M (1996) Sexual size dimorphism and determination of sex in Yellow-legged Gulls. *J Field Ornithol* 67: 534–541.
- Coulson JC, Duncan N, Thomas CS & Monaghan P (1981) An age-related difference in the bill depth of Herring Gulls *Larus argentatus*. *Ibis* 123: 499–502.
- Coulson JC, Thomas CS, Butterfield JEL, Duncan N, Monaghan P & Shedden C (1983) The use of head and bill length to sex live gulls Laridae. *Ibis* 125: 549–557.
- Croxall JP (1995) Sexual size dimorphism in seabirds. *Oikos* 73: 399–403.
- Evans DR, Hoopes ED & Griffin CR (1993) Discriminating the sex of Laughing Gulls by linear measurements. *J Field Ornithol* 64: 472–476.
- Fox GA, Cooper CR & Ryder JP (1981) Predicting the sex of Herring Gulls by using external measurements. *J Field Ornithol* 52: 1–9.
- Hokkaido Government (1999) *Government report for revival plan of community with seabirds* (in Japanese). Sapporo.
- Hokkaido Government (2000) *Government report for revival plan of community with seabirds* Hokkaido Government, Sapporo (in Japanese).
- Hunt GL Jr., Wingfield JC, Newman A & Farner DS (1980) Sex ratio of Western Gulls on Santa Barbara Island, California. *Auk* 97: 473–479.
- Ingolfsson A (1969) Sexual dimorphism of large Gulls (*Larus* spp.). *Auk* 86: 732–737.
- Jehl JR Jr. (1987) Geographic variation and evolution in the California Gull (*Larus californicus*). *Auk* 104: 421–428.
- Jones IL (1994) Mass changes of Least auklets *Aethia pusilla* during the breeding season: evidence for programmed loss of mass. *J Anim Ecol.* 63: 71–78.
- Lorentsen S-H & Røv N (1994) Sex determination of Antarctic Petrels *Thalassocia antarctica* by discriminant analysis of morphometric characters. *Polar Biol.* 14: 143–145.
- Monaghan P, Coulson JC, Duncan N, Furness RW, Shedden CB & Thomas C (1983) The geographical variation of the Herring Gull *Larus argentatus* within Britain and in northern Europe; a biometrical approach. *Ibis* 125: 412–417.

- Palomares LE, Arroyo B, Marchamalo J, Sainz JJ & Voslamber B (1997) Sex- and age-related biometric variation of black-headed Gulls *Larus ridibundus* in Western European populations. *Bird Study* 44: 310–317.
- Rodriguez EF, Pugsek BH & Diem KL (1996) A sexing technique for California Gulls breeding at Bamforth Lake, Wyoming. *J Field Ornithol* 67: 519–524.
- Ryder JP (1978) Sexing Ring-billed Gulls externally. *Bird-Banding* 49: 218–222.
- Schnell GD, Worthen GL & Douglas ME (1985) Morphometric assessment of sexual dimorphism in skeletal elements of California Gulls. *Condor* 87: 484–493.
- Shugart GW (1977) A method for externally sexing gulls. *Bird-Banding* 48: 118–121.
- Threlfall W & Jewer DD (1978) Notes on the standard body measurements of two populations of Herring Gulls (*Larus argentatus*). *Auk* 95: 749–753.
- Weimerskirch H & Jouventin P (1987) Population dynamics of the Wandering Albatross, *Diomedea exulans*, of the Crozet Islands: Cause and conservation of the population decline. *Oikos* 49: 315–322.