

Changes in Plasma Biochemistry and Body Mass During Incubation in the Yellow-legged Gull

CARLOS ALONSO-ALVAREZ^{1,3}, ALBERTO VELANDO², MIGUEL FERRER¹ AND JOSÉ A. R. VEIRA³

¹Department of Applied Biology, Estación Biológica de Doñana, CSIC, 41013 Sevilla, Spain
E-mail: alonso@ebd.csic.es

²Departamento de Ecoloxía e Bioloxía Animal, Universidade de Vigo, 36200 Vigo, Spain

³Departamento de Fisiología. Facultad de Veterinaria, Universidad de Santiago de Compostela. 27002 Lugo, Spain

Abstract.—The “Incubatory Reserves Constancy” hypothesis asserts that incubation could be a departure from breeding stress that allows for the maintenance or recovery of body reserves after laying effort (females) or territory defense (males) in those species with bi-parental incubation such as gulls. The plasma composition and body mass of incubating Yellow-legged Gulls (*Larus cachinnans*) were analyzed and related to the number of days after egg-laying. Female gulls showed an increase in uric acid and cholesterol levels, whereas males showed only an increase in uric acid values throughout this period. Moreover, females increased while males maintained their body masses. These results could reflect a recovery process after the laying effort supporting the Incubatory Reserves Constancy hypothesis in females. Uric acid and urea levels are positively correlated to body condition in Yellow-legged Gulls, which could be the result of a change in diet composition. This disagrees with recent findings on body composition in incubating gulls and could be related to variations in food availability among populations or years, and could reflect flexibility in the investment devoted by each sex. Received 4 December 2001, accepted 18 February 2002.

Key words.—Body-mass change, Incubatory Reserves Constancy Hypothesis, *Larus cachinnans*, plasma cholesterol, uric acid, urea, Yellow-legged Gull, triglycerides.

Waterbirds 25(2): 253-258, 2002

Reproduction requires a high energetic expenditure in birds (Drent and Daan 1980; Blem 1990). The breeding effort commonly results in a decrease of lipid and protein reserves (e.g., Ankney and MacInnes 1978; Alisauskas and Ankney 1994). However, the incubation period could be an exception to this general pattern in those species with bi-parental care, since the energy expenditure would be distributed between both sexes (Moreno 1989). Thus two opposing hypotheses may be proposed for changes in the energy reserves during incubation: the “Stress Hypothesis” would predict a pattern of gradual loss of reserves, because energetic costs are accumulated throughout the breeding season (Monaghan *et al.* 1989; Wendeln and Becker 1996), whereas “Incubatory Reserves Constancy Hypothesis” would suggest that the incubation period may be a departure from breeding stress that allows for the maintenance or recovery of body reserves (see Houston *et al.* 1983; Hario *et al.* 1991; also suggested by Moreno [1989] for total body mass).

Gulls are monogamous species with bi-parental incubation (Cramp and Simmons

1983) and both sexes incubate for a similar proportion of time (Drent 1970; Burger 1987; Alonso-Alvarez 2001; but see Butler and Janes-Butler 1983). Hario *et al.* (1991) and Mawhinney *et al.* (1999) analyzed body composition of Herring Gulls (*Larus argentatus*) and Great Black-backed Gulls (*Larus marinus*) respectively, which were culled throughout the breeding season by governmental agencies to control their numbers. Hario *et al.* (1991) found that both sexes gained mass and fat during incubation, whereas protein reserves remained stable. In contrast, Mawhinney *et al.* (1999) found that mass and lipid content decreased and protein increased in females, while mass and protein remained stable and fat increased in males. Therefore, the results of these studies suggest different energy expenditure for each sex.

In the present study, we analyzed four plasma biochemical parameters (uric acid, urea, triglycerides and cholesterol) and body mass in incubating Yellow-legged Gulls (*Larus cachinnans*) to determine changes in reserves of living individuals, thus trying to avoid sacrifice of birds. Plasma levels of uric

acid and urea reflect mobilization of protein reserves during food-shortage periods in birds (Cherel and Le Maho 1985; Boismenu *et al.* 1992; Alonso-Alvarez and Ferrer 2001). Plasma triglycerides value is related to body fat content (Bacon *et al.* 1989; Dabbert *et al.* 1997) and cholesterol level is a good indicator of body-mass loss from an individual's optimum in captive Yellow-legged Gulls (Alonso-Alvarez *et al.* 2002). However, few studies have analyzed changes in plasma chemistry throughout reproduction in free-living birds (Fairbrother *et al.* 1990; Cantos *et al.* 1994; Hollmén *et al.* 2001). Moreover, as far as we know, the plasma chemistry of breeder individuals has not been previously analyzed in any gull species.

The objectives of the present study were: (1) to determine whether the incubation period represents maintenance or recovery after the pre-laying effort in Yellow-legged Gull, as the Incubatory Reserves Constancy Hypothesis suggests, (2) to evaluate between-sexes differences in the patterns of change in body mass and plasma chemistry, and whether they are consistent with the results provided by Hario *et al.* (1991) or Mawhinney *et al.* (1999), and finally, (3) to report gulls' plasma biochemistry during the breeding period.

METHODS

The study was carried out on the Cies Islands (42°15'04"N, 8° 53'30"W), which are located in the outer area of the Ría de Viño (Spain) and are the breeding grounds for over 22,000 pairs of Yellow-legged Gulls. Birds were captured using nest-traps (Weaver and Kadlec 1970) during the incubation period (May 1998). We trapped 33 males and 43 females throughout the 27-31-d incubation period (Cramp and Simmons 1983). The beginning of incubation (following completion of clutch) was determined by visiting the colony every two d. In the last 10 d of incubation, 13 females and four males that had been trapped on the first 10 d were recaptured.

We weighed the birds with a spring balance to the nearest 5 g (Pesola). Two milliliters of blood were taken from each bird with a heparinized syringe from the brachial vein. Blood samples were collected between 11.00 h and 18.00 h to minimize variations caused by circadian rhythms (see Ferrer 1990). The blood was placed in tubes containing lithium heparin, which were kept on ice in cool containers (4°C) for a maximum period of eight h. Then, samples were centrifuged at 3,000 rpm for 10 min and plasma stored at -60°C until analysis.

Biochemical analyses were performed on a spectrophotometer (Hitachi-U2000, Tokyo, Japan). The deter-

minations were made in duplicate on each plasma sample following the methods and authors indicated in parentheses: uric acid (uricase method; Fossati *et al.* 1980), urea (urease method; Munan *et al.* 1978), triglycerides (peroxidase-coupled method; McGowan *et al.* 1983), and cholesterol (cholesterol esterase method; Alain *et al.* 1974). These parameters were analyzed using commercial reagents (Sigma-Aldrich Company, MO, USA). All samples were assayed in the same series. Uric acid: intra-assay CV = 1.9%, accuracy: ± 0.012 mmol/L. Urea: intra-assay Coefficient of Variation (CV) = 1.54%, accuracy: ± 0.33 mmol/L. Triglycerides: intra-assay CV = 1.5%; accuracy: ± 0.05 mmol/L. Cholesterol: intra-assay CV = 0.95%, accuracy: ± 0.08 mmol/L.

Data were tested for normality using the Shapiro-Wilk test. We used parametric tests because all variables were normally distributed. We tested between-sexes differences during first and last ten d of incubation by Student's *t*-test. Changes in plasma levels in the same individual between the first and the second sampling were tested by paired *t*-tests. Data of recaptured males were not used due to the small sample size. Relationships among variables were analyzed by Pearson correlation coefficients. In order to avoid pseudo-replication, recaptured males and females were not included in Pearson's correlations. An index of body condition (body mass corrected for size) was obtained as the residuals of the linear regression of body mass on cubed tarsus length of birds (males: $R^2 = 0.18$, $F_{1,31} = 6.74$, $P < 0.01$; females: $R^2 = 0.35$, $F_{1,41} = 22.08$, $P < 0.001$).

RESULTS

Female gulls showed lower urea and cholesterol concentrations than males during the first ten days of incubation (Table 1). There were no between-sexes differences in plasma levels during the middle ten days or during the last ten days of incubation (Table 1). Plasma concentration of uric acid in males ($t_{21} = 2.42$, $P < 0.05$) and uric acid and cholesterol in females ($t_{22} = 3.13$, $P < 0.01$; $t_{22} = 2.93$, $P < 0.01$; respectively) showed significant increase between the first and the last 10-d periods (see Table 1). Moreover, the 13 females recaptured in the last days of incubation showed a significant increase in uric acid and cholesterol values (Table 2). Plasma variables and body condition index were not correlated with the laying date (n.s. in all cases and in both sexes).

In males, only uric acid showed a significant correlation with the number of days spent in incubation before blood extraction (Table 3). In females, uric acid, urea, cholesterol and body condition index were positively correlated with the day of incubation (Table 3). In addition, body mass was positively correlated with the day of incubation

Table 1. Concentrations of four plasma chemicals in the Yellow-legged Gull during the first and last days of incubation period. Values are in mmol/L and are given as the mean ± 1 SE.

Parameter	First 10 days ^a				Middle 10 days ^b				Last 10 days ^c							
	Males		Females		Males		Females		Males		Females		P			
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	t ₁₉	P		
Uric acid	0.46 ± 0.05		0.57 ± 0.10		1.09		0.65 ± 0.1		0.63 ± 0.11		0.12		0.93 ± 0.18		1.10	n.s.
Urea	1.01 ± 0.13		0.57 ± 0.12		2.55		0.98 ± 0.09		0.82 ± 0.08		1.25		0.94 ± 0.16		0.06	n.s.
Triglycerides	1.06 ± 0.17		1.31 ± 0.11		1.26		0.92 ± 0.09		1.40 ± 0.2		1.73		1.41 ± 0.73		1.82	n.s.
Cholesterol	7.35 ± 0.32		5.82 ± 0.40		2.81		7.24 ± 0.42		6.94 ± 0.32		0.57		7.82 ± 0.35		0.41	n.s.

Student's t-test for independent samples.

^a11 males and 15 females.

^b10 males and 19 females.

^c12 males and 9 females.

in females ($r_{41} = 0.33, P < 0.05$), but not in males ($r_{31} = 0.01, n.s.$). Finally, body condition index was positively correlated with uric acid ($r_{41} = 0.54, P < 0.001$) and urea ($r_{41} = 0.37, P < 0.05$) in females, and with uric acid in males ($r_{31} = 0.43, P < 0.05$).

DISCUSSION

Breeding produces marked changes in the body mass of gulls (Coulson *et al.* 1983) and, as shown here, also in plasma composition during incubation. Body condition index, body mass and plasma biochemistry of Yellow-legged Gulls did not indicate a decrease in body reserves throughout this period. Hence, these results suggest that the "Stress Hypothesis" may be rejected in this species.

Nor do our findings support another hypothesis proposed to explain mass decrease during breeding, which is known as the "Programmed Anorexia Hypothesis" (Norberg 1981; Gaston and Jones 1989). It suggests that lipid storage levels may be minimized to maintain a low wing loading, thus increasing the flight efficiency. This would be an adaptive trait in seabirds since improves the foraging efficiency during the chick rearing period (i.e., Gaston and Jones 1989; Croll *et al.* 1991).

However, the results found in the present study may support the "Incubatory Reserves Constancy" hypothesis. Furthermore, male and female gulls differed in their patterns, such as Mawhinney *et al.* (1999) found. We detected an increase in body mass in females, but not in males during incubation. This might be related to between-sexes differences in energy expenditure before incubation, thus females recovering their previous state. In fact, females invest in egg production, which appreciably increases their energetic requirements. Unfortunately, we could not trap birds in the colony during the pre-laying period to test a recovery pattern. However, there were significant between-sexes differences in plasma composition during the first incubation days (see Table 1).

Plasma urea and uric acid are waste products of protein catabolism (Griminger and Scanes 1986). In birds, high levels have been associated with starvation and body mass

Table 2. Values of four plasma chemicals (mmol/L) and body mass (g) of 13 female Yellow-legged Gulls captured in the first and last ten days of incubation. Values are given as mean \pm 1 SE.

	First 10 days	Last 10 days	Change	t_{12}	P
Uric acid					<0.05
Urea					n.s.
Triglycerides					n.s.
Cholesterol					<0.05
Body mass					n.s.

Student's t-paired test.

loss, because muscular protein is used as energy source during the last phase of fasting (Cherel and Le Maho 1985; Boismenu *et al.* 1992; Castellini and Rea 1992). In the same way, plasma urea has been used as a nutritional condition index (Ferrer 1992, 1993). The protein is a limiting factor during albumen production among bird species (Williams 1996; Ramsay and Houston 1997), including gulls (Herring Gull; Houston *et al.* 1983). As female gulls differentially suffered the costs of egg production, we should expect high nitrogen values at the start of incubation, decreasing throughout this period. Surprisingly, we found the opposite.

In a recent study, Hollmén *et al.* (2001) found an increase of uric acid plasma levels in female Common Eiders (*Somateria mollissima*) during incubation. The Common Eider has uni-parental incubation. Female Eiders rarely leave their nest to feed during this period (26 d), relying energetically on their body reserves to lost approximately 30% of their body masses (Parker and Holm 1990). Hollmén *et al.* (2001) related the uric acid increase to mobilization of protein reserves

and body-mass loss. However, our results suggest that in other circumstances this relationship may be reversed. Both urea and uric acid were positively correlated with body mass corrected for body size in Yellow-legged Gull. Hence, such relationships must be carefully considered. A possible explanation might derive from changes in diet.

A rise in the proportion of protein in the diet increased the plasma level of these nitrogen metabolites in chickens and pigeons (Okumura and Tasaki 1969; Featherston 1969; Lumeij and Bruijne 1985). Therefore, our results relate to diet differences, allowing female gulls to increase their protein reserves. The studied birds often fed on refuse tips (rubbish present in the 40% of pellets [Munilla 1997a]), and showed marked modifications in diet throughout the breeding period, as the occurrence of crabs in regurgitations increases significantly during incubation (Munilla 1997b). An increase in the proportion of crustaceans has also found in the diet of Kelp Gulls (*Larus dominicanus*) in the same period (Bertelotti and Yorio 1999).

On the other hand, triglyceride values did not change during incubation in either sex, indicating stable lipid reserves, since the amount of body fat is positively correlated with plasma triglycerides values (Bacon *et al.* 1989; Dabbert *et al.* 1997). Furthermore, cholesterol is lower in females during first ten days, increasing during the rest of incubation period. In Yellow-legged Gulls, the plasma concentration of this lipid is negatively correlated with the individual proportion of body-mass loss, thus providing a reliable index of the body condition (Alonso-Alvarez *et al.* 2002). Incubating gulls with high cholesterol levels would be relatively heavy birds. The

Table 3. Pearson correlation between the day of incubation and four plasma parameters and residuals of body mass on body size (body condition).

Parameter	Males ^a		Females ^b	
	r_{31}	P	r_{41}	P
Uric acid	0.341	<0.05	0.538	<0.001
Urea	-0.061	n.s.	0.322	<0.05
Triglycerides	0.238	n.s.	0.228	n.s.
Cholesterol	0.229	n.s.	0.399	<0.01
Body condition	0.099	n.s.	0.389	<0.05

cholesterol increase bears out the presence of a recovery pattern in female birds.

In summary, the present results support the hypothesis that body mass and reserves maintain or recover their levels during incubation in those species with bi-parental care. However, whereas females increased body mass, and probably changed diet composition in order to recuperate the protein expenditure during laying effort, males maintained body mass, lipids and protein metabolites throughout incubation. These results are not consistent with the results of Hario *et al.* (1991) on Herring Gulls or Mawhinney *et al.* (1999) on Great Black-backed Gulls. The differences between the two studies were explained by differences in the energy expenditure of the sexes during incubation (Mawhinney *et al.* 1999). Female Great Black-backed Gulls spent more time on the nest than males, thus reducing her feeding time (Butler and Janes-Butler 1983). Alternatively, the disagreement between Hario *et al.* (1991), Mawhinney *et al.* (1999) and the present paper might be related to differences in food availability (Kitaysky *et al.* 1999). Differences in food abundance produced a readjustment of each sex role during incubation in Western Gulls (*Larus occidentalis*; Pierotti 1981). The plasma chemistry during the incubation period should be analyzed under different food-availability situations, which could be easily manipulated by means of an artificial food supply. Moreover, the relationship between food composition and plasma biochemistry in wild birds should be examined.

ACKNOWLEDGMENTS

We are grateful to the staff of Islas Cies Natural Park. We thank to J. L. Tella, Bertellotti, M. and V. Penteriani for their valuable comments. The capture and sampling of birds were conducted under legal permission from the wildlife authorities (Xunta de Galicia), avoiding unnecessary damages to the birds.

LITERATURE CITED

- Alisaukas, R. T. and C. D. Ankney. 1994. Nutrition of breeding female Ruddy Ducks: the role of nutrient reserves. *Condor* 96: 878-897.
- Allain, C. A., L. S. Poon, C. S. G. Chan, W. Richmond and P. C. Fu. 1974. Enzymatic determination of total serum cholesterol. *Clinical Chemistry* 20: 470-478.
- Alonso-Alvarez, C. 2001. Effects of testosterone implants on pair behaviour during incubation in the Yellow-legged Gull *Larus cachinnans*. *Journal of Avian Biology* 32: 326-332.
- Alonso-Alvarez, C. and M. Ferrer. 2001. Biochemical study about fasting, subfeeding and recovery processes in Yellow-legged Gulls. *Physiological and Biochemical Zoology* 74: 703-713.
- Alonso-Alvarez, C., M. Ferrer and A. Velando. 2002. The plasmatic index of body condition in Yellow-legged Gulls (*Larus cachinnans*): a food-controlled experiment. *Ibis* 144: 147-150.
- Ankney, C. D. and C. D. MacInnes. 1978. Nutrient reserves and reproductive performance of female Lesser Snow Geese. *Auk* 95: 459-471.
- Bacon, W. L., K. E. Nestor and E. C. Naber. 1989. Prediction of carcass composition of turkeys by blood lipids. *Poultry Science* 68: 1282-1288.
- Bertellotti, M. and P. Yorio. 1999. Spatial and temporal patterns in the diet of the Kelp Gull in Patagonia. *Condor* 101: 790-798.
- Blem, C. R. 1990. Avian energy storage. Pages 59-113 in *Current Ornithology*, Vol. 7 (D. M. Power, Ed.). Plenum Press, New York.
- Boismenu, C., G. Gauthier and J. Larrochelle. 1992. Physiology of prolonged fasting in Greater Snow Geese (*Chen caerulescens atlantica*). *Auk* 109: 511-521.
- Burger, J. 1987. Selection for equitability in some aspects of reproductive investment in Herring Gulls *Larus argentatus*. *Ornis Scandinavica* 18: 17-23.
- Butler, R. G. and S. Janes-Butler. 1983. Sexual differences in the behaviour of the adult Great Black-backed Gulls (*Larus marinus*) during the pre-and-post hatch periods. *Auk* 100: 63-75.
- Cantos, F. J., A. L. Alonso-Gómez and M. J. Delgado. 1994. Seasonal changes in fat and protein reserves of the Black-headed Gull, *Larus ridibundus*, in relation to migration. *Comparative Biochemistry and Physiology* 108A: 117-122.
- Castellini, M. A. and L. D. Rea. 1992. The biochemistry of natural fasting at its limits. *Experientia* 48: 575-582.
- Cherel, Y. and Y. Le Maho. 1985. Five months of fasting in King Penguin chicks: body mass loss and fuel metabolism. *American Journal of Physiology* 249: R387-392.
- Coulson, J. C., P. Monaghan, J. Butterfield, N. Duncan, C. Thomas and C. Shedden. 1983. Seasonal changes in the Herring Gull in Britain: weight, moult and mortality. *Ardea* 71: 235-244.
- Cramp, S. and K. E. L. Simmons. 1983. *Handbook of the birds of Europe, The Middle East and North Africa. The birds of Western Palearctic. Vol. III. Waders to Gulls.* Oxford University Press, Oxford.
- Croll, D. A., A. J. Gaston and D. G. Noble. 1991. Adaptive loss of mass in thick-billed murre. *Condor* 93: 496-502.
- Dabbert, C. B., T. E. Martin and K. C. Powell. 1997. Use of body measurements and serum metabolites to estimate the nutritional status of Mallards wintering in the Mississippi Alluvial Valley, USA. *Journal of Wildlife Diseases* 33: 57-63.
- Drent, R. H. 1970. Functional aspects of incubation in the Herring Gull (*Larus argentatus*). *Behaviour* 17: S1-S132.
- Drent, R. H. and S. Daan. 1980. The prudent parent: energetic adjustments in avian breeding. *Ardea* 68: 225-252.
- Fairbrother, A., M. A. Craig, K. Walker and D. O'Loughlin. 1990. Changes in Mallard (*Anas platyrhynchos*) serum chemistry due to age, sex and reproductive condition. *Journal of Wildlife Diseases* 26: 67-77.

- Featherston, W. R. 1969. Nitrogenous metabolites in the plasma of chicks adapted to high protein diets. *Poultry Science* 48: 646-651.
- Ferrer, M. 1990. Hematological studies in birds. *Condor* 92: 1085-1086.
- Ferrer, M. 1992. Natal dispersal in relation to nutritional condition in Spanish Imperial Eagles. *Ornis Scandinavica* 23: 104-107.
- Ferrer, M. 1993. Ontogeny of dispersal distances in young Spanish Imperial Eagles. *Behavioral Ecology and Sociobiology* 32: 259-263.
- Fossati, P., L. Prencipe and G. Berti. 1980. Use of 3,5-dichloro-2-hydroxybenzene-sulfonic acid/4-aminophenazone chromogenic system in direct enzymatic assay of uric acid in serum and urine. *Clinical Chemistry* 26: 227-231.
- Gaston, A. J. and I. L. Jones. 1989. The relative importance of stress and programmed anorexia in determining mass loss by incubating ancient murrelets. *Auk* 106: 653-658.
- Griminger, P. and C. G. Scanes. 1986. Protein metabolism. Pages 326-344 in *Avian Physiology* (P. D. Sturkie, Ed.). Springer-Verlag, New York.
- Hario, M., M. Kilpi and K. Selin. 1991. Parental investment by sexes in Herring Gull: the use of energy reserves during the early season. *Ornis Scandinavica* 22: 308-312.
- Hollmén, T., J. C. Franson, S. Sankari, M. Kilpi and K. Lindström. 2001. Use of serum biochemistry to evaluate nutritional status and health of incubating Common Eiders (*Somateria mollissima*) in Finland. *Physiological and Biochemical Zoology* 74: 333-342.
- Houston, D. C., P. J. Jones and R. M. Sibly. 1983. The effect of female body condition on egg laying in Lesser Black-backed Gulls *Larus fuscus*. *Journal of Zoology, London* 200: 509-520.
- Kitaysky, A. S., J. C. Wingfield and J. F. Piatt. 1999. Dynamics of food availability, body condition and physiological stress in breeding Black-legged Kittiwakes. *Functional Ecology* 13: 577-584.
- Lumeij, J. T. and J. J. Bruijne. 1985. Blood chemistry reference values in Racing Pigeons (*Columba livia domestica*). *Avian Pathology* 14: 401-408.
- Mawhinney, K., A. W. Diamond and F. P. Kehoe. 1999. The use of energy, fat, and protein reserves by breeding Great Black-backed Gulls. *Canadian Journal of Zoology* 77: 1459-1464.
- McGowan, M. W., J. D. Artiss, D. R. Strandberg and B. Zak. 1983. A peroxidase-coupled method for the colorimetric determination of serum triglycerides. *Clinical Chemistry* 29: 538-542.
- Monaghan, P., J. D. Uttley, M. D. Burns, C. Thaine and J. Blackwood. 1989. The relationship between the food supply, reproductive effort and breeding success in Arctic Terns *Sterna paradisaea*. *Journal of Animal Ecology* 58: 261-274.
- Moreno, J. 1989. Strategies of mass change in breeding birds. *Biological Journal of the Linnean Society* 37: 297-310.
- Munan, L., A. Kelly, C. PetitClerc and B. Billon. 1978. Associations with body weight of selected chemical constituents in blood: epidemiologic data. *Clinical Chemistry* 24: 772-777.
- Munilla, I. 1997a. Estudio de la Población y la Ecología Trófica de la Gaviota Patiamarilla, *Larus cachinnans* Pallas, en Galicia. Unpublished Ph.D. Thesis. University of Santiago de Compostela, Santiago de Compostela.
- Munilla, I. 1997b. Henslow's Swimming Crab (*Polydora henslowii*) as an important food for Yellow-legged Gulls (*Larus cachinnans*) in NW Spain. *I.C.E.S. Journal of Marine Sciences* 54: 631-634.
- Norberg, R. A. 1981. Temporary weight decrease in breeding birds may result in more fledged young. *American Naturalist* 118: 838-850.
- Okumura, J. and I. Tasaki. 1969. Effect of fasting, re-feeding and dietary protein level on uric acid and ammonia content of blood, liver and kidney in chickens. *Journal of Nutrition* 97: 316-320.
- Parker, H. and H. Holm. 1990. Patterns of nutrient and energy expenditure in female Common Eiders nesting in the high Arctic. *Auk* 107: 660-668.
- Pierotti, R. 1981. Male and female parental roles in the Western Gull under different environment conditions. *Auk* 98: 532-549.
- Ramsay, S. L. and D. C. Houston. 1997. Nutritional constraints on egg production in the Blue Tit: a supplementary feeding study. *Journal of Animal Ecology* 66: 649-657.
- Weaver, D. K. and J. A. Kadlec. 1970. A method for trapping breeding adult gulls. *Bird-Banding* 41: 28-31.
- Wendeln, H. and P. H. Becker. 1996. Body mass change in breeding Common Terns *Sterna hirundo*. *Bird Study* 43: 85-95.
- Williams, T. D. 1996. Variation in reproductive effort in female Zebra Finches (*Taeniopygia guttata*) in relation to nutrient-specific dietary supplements during egg laying. *Physiological Zoology* 69: 1255-1275.